

# PHARMACEUTICAL COMPOSITIONS COMPRISING AMLODIPINE MALEATE

This application is a continuing application under 35 U.S.C. § 120 of prior co-  
pending U.S. patent application Serial No. 09/809,346, filed March 16, 2001, the  
entire contents of which are incorporated herein by reference. Further, this  
application claims the benefit of priority under 35 U.S.C. § 119(e) from U.S.  
provisional application 60/258,562, filed December 29, 2000, the entire contents of  
which are incorporated herein by reference.


## BACKGROUND OF THE INVENTION

### 1. Field of the Invention

The present invention relates to pharmaceutical compositions comprising  
amlodipine maleate and to processes for making the same.

### 2. Description of the Related Arts

Calcium channel blockers are useful in treating a variety of cardiac conditions,  
primarily angina and hypertension. EP 089 167 and corresponding US 4,572,909  
disclose a class of substituted dihydropyridine derivatives as being useful calcium  
channel blockers. These patents identify that one of the most preferred compounds is  
2-[(2-aminoethoxy)methyl]-4-(2-chlorophenyl)-3-ethoxycarbonyl-5-  
methoxycarbonyl-6-methyl-1,4-dihydropyridine. This compound, which is now  
commonly known as amlodipine, has the following formula:



Examples 8, 11, 12, and 22 of EP 089 167 show the synthesis of amlodipine in  
the maleate salt form. While a variety of acid addition salts are taught to be suitable,  
the maleate salt is identified as the most preferred acid addition salt.

Subsequently, EP 244 944 and corresponding US 4,879,303 were issued  
directed to the besylate (or benzene sulfonate) salt of amlodipine. The besylate salt is  
stated to provide certain advantages over the known salts including good formulating

properties. Indeed, amlodipine besylate, and not amlodipine maleate, has been developed into a commercial prescription pharmaceutical by Pfizer and is sold in the U.S. under the trade name NORVASC.

A review of the available portions of the NORVASC (amlodipine besylate) New Drug Application (NDA) filed with the U.S. Food & Drug Administration by Pfizer reveals that a switch was made during development from the original amlodipine maleate to the amlodipine besylate. (See "Review of Original NDA" for NDA# 19-787 of 10.10.1990, obtainable from FDA under Freedom of Information Act ). The reasons for the switch were apparently tableting and stability problems. However, the precise stability and tableting problems/issues/causes are not publicly disclosed in the information available from the FDA. Interestingly, clinical studies in the NDA include the use of the maleate salt form and the besylate salt form, the two salt forms being therapeutically equivalent (bioequivalent). However, in these studies amlodipine maleate was always used in a capsule or solution dosage form, not a tablet dosage form.

#### SUMMARY OF THE INVENTION

The present invention relates to the discovery of stable amlodipine maleate pharmaceutical compositions. A first aspect of the invention relates to a pharmaceutical composition comprising an effective amount of amlodipine maleate and at least one pharmaceutically acceptable excipient wherein the composition has a pH within the range of 5.5 - 7.0. The composition is preferably a solid dosage form such as a tablet or capsule. The invention also relates to methods of making such a composition as well as to the use of such compositions in treating or preventing angina or hypertension.


Another aspect of the invention relates to a process which comprises mixing solid particles of amlodipine maleate having an average particle size of at least 20 microns with one or more pharmaceutically acceptable excipients to form a mixture. Preferably the mixture has a pH of from 5.5 to 7.0. The mixture can be compressed into tablets or filled into capsules to form a solid dosage form.

#### DETAILED DESCRIPTION OF THE INVENTION

Stability is an important aspect of a pharmaceutical composition. The present invention is based on the discovery that the prior art stability problems associated


with amlodipine maleate can be overcome primarily by controlling the pH of the composition to be within the range of about 5.5 to 7.0, preferably of about 6.0-7.0. Within this range the potential degradation reactions are minimized. In particular, the formation of the following degradation product, referred to herein as amlodipine aspartate, is diminished or prevented in this pH range.

#### Amlodipine Aspartate



Amlodipine aspartate is formed by a Michael addition reaction between amlodipine and maleic acid. Because amlodipine and maleic acid are in intimate contact with each other in the amlodipine maleate salt, the chances of the addition reaction occurring increase with time; hence raising a stability issue. By controlling the pH, the addition is significantly slowed or prevented altogether. Thus, it has now been discovered that pH levels above 7.0 tend to encourage the degradation of amlodipine maleate into amlodipine aspartate. Below a pH of about 5.5, other degradation reactions tend to be encouraged, including the pyridine analogue of amlodipine having the following structure:

#### “Amlo-pyridine”



Preferably the pH of the stabilized composition of our invention is in the range of about 6.0 to 7.0 and more typically from about 6.1 or 6.2 to 6.8. For solid compositions, the pH is determined by forming a slurry of the material with water (demineralized water) and measuring the pH of the slurry, as is understood by

workers skilled in the art regarding the pH of a solid composition. The concentration of the composition in the slurry is 20 wt%. The pH is measured by any standard technique.

The pharmaceutical compositions of the present invention comprise a pharmaceutically effective amount of amlodipine maleate and at least one pharmaceutically acceptable excipient. Preferably the stability of the composition is such that after three months, more preferably after six months, in an environment controlled room at 40°C/75% RH, it exhibits a loss of amlodipine (or, accordingly, an increase in impurities content) of less than 10 %, preferably less than 5 %, and more preferably less than 1%. Alternatively, the amlodipine maleate pharmaceutical compositions of the present invention preferably exhibit a storage stability equivalent to or superior to amlodipine besylate compositions. For example, the loss of amlodipine during storage to degradation reactions is equivalent to (+/- 10%) or less than the amlodipine loss in amlodipine besylate compositions, especially the commercial product.

The form of the amlodipine maleate is not particularly limited and includes anhydrides, solvates, hydrates and partial hydrates as well as crystalline and amorphous forms. Further, the ratio of amlodipine to maleate can be varied and specifically includes the more common and prior art form of 1:1 as well as a novel 2:1 form described commonly owned co-pending U.S. patent application serial No. 09/809,356, filed on March 16, 2001, and entitled "Amlodipine Hemimaleate," the entire contents of which are incorporated herein by reference.

The amount of amlodipine is not particularly limited and includes any amount that provides a pharmaceutical effect. In particular, amlodipine maleate can be used to treat or prevent hypertension or angina by administering an effective amount to a patient in need thereof. The specific form of angina is not particularly limited and specifically includes chronic stable angina pectoris and vasospastic angina (Prinzmetal's angina). The compound can be administered by any suitable route including orally or parenterally depending on the dosage form. The "patients" intended to be treated include human and non-human animals especially non-human mammals. Generally the amount of amlodipine maleate in a unit dose is from 1 to 100 mg, more typically from 1 to 25 mg, and preferably about 1, 1.25, 2.5, 5 or 10 mg (expressed in terms of the free base).



acidic. For example, amlodipine maleate has a pH of about 4.8 as a saturated aqueous solution. Thus using excipients that are pH inert; i.e., they have little to no effect on pH, generally results in a non-alkaline pharmaceutical composition because the amlodipine maleate essentially acts as its own adjusting agent. An example of a pH-inert excipient is microcrystalline cellulose. A composition comprising amlodipine maleate and microcrystalline cellulose generally exhibits a pH of about 6. Commercially available tablets comprising amlodipine besylate and sold under brand name Norvasc exhibit a pH typically between 7.05-7.35 (again measured as a 20 wt% slurry).

Excipients having a pH effect can also be used. The pH of these excipients must be taken into account in developing the pharmaceutical composition so that the overall pH of the pharmaceutical composition falls within the range of about 5.5 to 7.0.

For example, the commercially available/pharmaceutically acceptable calcium phosphates are generally alkaline; i.e. pH greater than 7 when measured as described above in a 20% slurry. For instance, DI-TAB, a commercially available dibasic calcium phosphate dihydrate, is reported as having a pH of about 7.4. Nonetheless some forms and grades of calcium phosphate are acidic or neutral pH. This lower pH can be due to the species of calcium phosphate as well as the treatment during processing of the material, such as in removing impurities/washing. For example, dibasic calcium phosphate anhydrate is generally considered to have a pH of about 7.3 whereas A-Tab <sup>TM</sup> (Rhodia), also a dibasic calcium phosphate anhydrate, has a pH of about 5.0. Further examples of commercially available non-alkaline calcium phosphates include DiCAFOS A (Budenheim) having a pH of about 7 (10 % slurry) and Fujicalin SG (Fuji) having a pH from 6.1-7.2 (5 % slurry). By using a non-alkaline calcium phosphate as an excipient, a pharmaceutical composition meeting the desired pH can be attained. Alternatively, a blend of calcium phosphates, some at pH above 7 and some a pH below 7 can be used to achieve the desired pH of the composition.

Instead of, or in addition to a non-alkaline calcium phosphate, other acidic excipients can be used by themselves or to counter balance an alkaline excipient. An example of such an acidic excipient is the disintegrant Explotab(TM) of Penwest, which is a cross-linked, low substituted sodium starch glycollate. Further, pH adjusting agents can also be used to attained the desired pH. These agents include

pharmaceutically acceptable acids such as maleic acid, citric acid or ascorbic acid (the second two have utility also to act as antioxidants) and pharmaceutically acceptable bases such as calcium oxide or magnesium oxide. Salts of weak acids and/or weak bases are also suitable pH adjustors as they act as buffers imparting pH to lower or  
5 higher values according to the chemical nature of their components.

The pharmaceutical compositions of the present invention are not particularly limited in terms of form or route of administration. Oral dosage forms as well as parenteral dosage forms are included. The composition can be in the form of a liquid, solid, or suspension. Preferably, the pharmaceutical composition is a solid dosage  
10 form such as a tablet, capsule or sachet intended for oral administration.

Preferred solid dosage forms contain as a major excipient microcrystalline cellulose, a calcium phosphate, especially calcium hydrogenphosphate, or a combination thereof. The sum of the other excipients, if any, is generally less than 25 wt%, more usually less than 10 wt% and in some cases less than 5 wt% of the total  
15 pharmaceutical composition. Preferred other excipients are a disintegrant such as sodium starch glycolate and/or a lubricant such as magnesium stearate and/or talc.

For example, a pharmaceutical composition comprising amlodipine maleate and microcrystalline cellulose as the only excipient has been shown to provide good stability against the formation of impurities.

20 The pharmaceutical compositions of the present invention can be made by techniques generally known in the art. In general, the amlodipine maleate is mixed with one or more excipients to form a blend. The mixing can be carried out wet or dry (i.e. using or not using a solvent or a liquid diluent in the process) and can involve granulating, slugging, or blending of powders. A dry process is however preferred.  
25 The blend, after optional further processing, can be compressed into tablets or filled into capsules such as gelatin capsules. Typically the amlodipine maleate to be mixed is in the form of particles. The storage stability of the pharmaceutical composition of the present invention is enhanced, in general, by using larger particle sizes. Preferably the average particle size of the amlodipine maleate is at least 20 microns,  
30 more preferably at least 100 microns, and in some embodiments at least 300 microns. If an adjustment of the pH of the composition is needed, it preferably is adjusted before processing into the final form such as a tablet or capsule.

For example, tablets made of the present invention can be prepared, e.g. by a wet granulation of a mixture of amlodipine maleate and a solid carrier/diluent such as

calcium phosphate of the proper grade, with the aid of a granulating solvent such as water or ethanol; drying the wet granulate; sieving the granulate; blending with sodium starch glycollate and magnesium stearate and compressing the tablet mixture into tablets. The control of pH and/or adjusting the pH value should be advantageously included before blending with magnesium stearate. In this example both the granulating step and the blending step are considered as "mixing" steps since amlodipine maleate and an excipient are mixed.

Yet another suitable process comprises direct compression of the blend of amlodipine maleate and excipient(s). In this process, the ingredients are blended together to form a compressible blend composition that is subsequently compressed into a tablet. A blend that comprises amlodipine maleate, microcrystalline cellulose and/or calcium phosphate, and optionally sodium starch glycollate and/or magnesium stearate may be useful in forming a tablet by direct compression. For example, a blend comprising amlodipine maleate, calcium hydrogen phosphate, microcrystalline cellulose and sodium starch glycollate having a pH of 5.5 to 7.0 can be blended together, re-blended with magnesium stearate and pressurized to form a stable tablet.

The tablets of the present invention comprising amlodipine maleate, microcrystalline cellulose, sodium starch glycollate and magnesium stearate, optionally with addition of calcium hydrogenphosphate, does not suffer from the problem of stickiness to the tablet punch as reported in prior art regarding other amlodipine formulations (see the above cited EP 244 944). Accordingly, the composition of the present invention may be produced in industrial scale without technological problems.

Tablets may be covered with a suitable coating. For example, the coating can be a moisture or light barrier to help with storage stability or a sustained or delayed release coating composition as are well known in the art.

Alternative dosage forms are capsules, both soft and hard capsules. The stabilized amlodipine maleate composition of the invention as described above is filled into capsules by techniques known in the art, in amounts comprising the desired therapeutical dose of amlodipine.

Suitable package material for packing the pharmaceutical dosage forms are plastic or glass containers and blisters. Particularly blisters made from non-permeable materials (high density polyethylene or aluminium) are advantageous as they may

contribute to decreasing the rate of formation of degradation impurities, namely the amlo-pyridine impurity, during storage.

The pharmaceutical compositions of the present invention are used in treating or preventing angina or hypertension, as previously defined, by administering an effective amount of the pharmaceutical composition to a patient in need thereof. Typically the pharmaceutical composition is a unit dose. Individual unit dose compositions generally contain from 1 to 100 mg of amlodipine maleate, more usually from 1 to 25 mg. Preferable are unit doses comprising amlodipine maleate in an equivalent of 1.25, 2.5, 5 or 10 mg of amlodipine, such as tablets or capsules for oral administration. The pharmaceutical composition is administered from 1 to 3 times daily, preferably once a day. The above compositions are useful also in reducing of heart failure symptoms, improving systolic left ventricular function and increasing exercise capacity in patients with ischaemic LVD and heart failure without current angina.

Amlodipine maleate compositions of our invention may be also used in medical applications in combination with other antihypertensive and/or antianginal agents, for instance with ACE- inhibitors such as benazepril. The combination may be realized in a form of single combination preparation, e.g. a capsule containing amlodipine maleate and benazepril hydrochloride, or by separate administration of drugs containing the above agents. Similarly, amlodipine maleate may also be combined with HMG-CoA reductase inhibitors such as lovastatin, simvastatin, atorvastatin as well as other statins.

Accordingly, the present invention further provides a method for treating and/or preventing any one or more of angina and hypertension by administering a pharmaceutical composition of our invention comprising an effective and/or prophylactic amount of amlodipine maleate to a sufferer in need thereof.

The present invention also provides the use of the composition of the invention in the manufacture of a medicament for treating and/or preventing any one or more of the Disorders.

## Examples

### **EXAMPLE 1 Amlodipine maleate tablets based on calcium phosphate**

#### **excipient**

5

a) Tablet compositions comprising calcium phosphate of different pH

Batch number	(A)	(B)	(C)	(D)	(E)	(F)
Equivalent of amlodipine	2.5 mg	10 mg	2.5 mg	10 mg	2.5 mg	10 mg
Amlodipine maleate	3.21 mg	12.8 mg	3.21 mg	12.8 mg	3.21 mg	12.8 mg
Calcium hydrogen phosphate anhydrous: - Di CAFOS A - A-TAB - Fujicalin	31.5 mg	126.0 mg	31.5 mg	126.0 mg	31.5 mg	126.0 mg
Microcrystalline cellulose	62.05 mg	248.1 mg	62.05 mg	248.1 mg	62.05 mg	248.1 mg
Sodium starch glycollate	2.0 mg	8.0 mg	2.0 mg	8.0 mg	2.0 mg	8.0 mg
Magnesium Stearate	1.0 mg	4.0 mg	1.0 mg	4.0 mg	1.0 mg	4.0 mg
Total	99.76 mg	398.9 mg	99.76 mg	398.9 mg	99.76 mg	398.9 mg

#### **Types of anhydrous calcium hydrogenphosphate used**

10

Batch no.	Final pH of 20% (m/V) slurry of tabs	Type of CaHPO <sub>4</sub>	Supplier of CaHPO <sub>4</sub>	pH of 5% (m/V) slurry of CaHPO <sub>4</sub>	pH of 20% (m/V) slurry of CaHPO <sub>4</sub>
(A), (B)	6.13, 6.19	Di CAFOS A	Budenheim	7.29	6.69
(E), (F)	5.74, 5.74	Fujicalin	Fuji	6.12	5.62
(C), (D)	5.53, 5.54	A-TAB	Rhone-Poulenc	6.03	5.25

b) Tablet compositions comprising amlodipine maleate of different particle sizes

Batch number	(G)	(H)	(I)
Equivalent of amlodipine base	2.5 mg	10 mg	2.5 mg
Amlodipine maleate, milled	3.21 mg	12.8 mg	-
Amlodipine maleate	-	-	3.21 mg
Calciumhydrogenphosphate anhydrous ( pH 6.7 )	31.5 mg	126.0 mg	31.5 mg
Microcrystalline cellulose	62.05 mg	248.1 mg	62.05 mg

Sodium starch glycollate	2.0 mg	8.0 mg	2.0 mg
Magnesium Stearate	1.0 mg	4.0 mg	1.0 mg
Total	99.76 mg	398.9 mg	99.76 mg

Particle size of the amlodipine maleate substance used for production of batches A – F was measured by laser diffraction and it was proven that less than 90% of particles is smaller than 204 microns and 50% of particles is smaller than 80 microns.

This amlodipine maleate was milled to a particle size of 10-20 microns and used for producing batches (G) and (H). Alternately, another batch of amlodipine maleate with particle sizes 90% less than 11 microns and 50% less than 6 microns was used for producing batch (I).

10

#### c) Manufacturing procedures

##### **Batches (A) – (F) and (I) have been manufactured as follows:**

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- 15 - All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 2.5 mg and / or 10 mg tablets have been compressed using a Korsch EK0 excenter
- 20 press.

##### **Batches (G) and (H) have been manufactured as follows:**

- Amlodipine maleate was milled
- The other excipients have been sieved through a 850 µm screen.
- 25 - All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 2.5 mg / 10 mg tablets have been compressed on Korsch EK0 excenter press.

30

### **Example 2 Tablets comprising microcrystalline cellulose**

#### **a) Composition**

Batch number	(J)	(K)
Equivalent of amlodipine base	2.5 mg	10 mg
Amlodipine maleate	3.21 mg	12.8 mg
Microcrystalline cellulose	75.55 mg	302.1 mg
Predried potato starch	20.0 mg	80.0 mg
Magnesium Stearate	0.5 mg	2.0 mg
Talc	0.5 mg	2.0 mg
Total	99.76 mg	398.9 mg

#### **Manufacturing process**

- 5            -        Amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate and talc have been mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- 10           -        Magnesium stearate and talc were added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 2.5 mg and 10 mg tablets have been compressed using a Korsch EK0 excenter press.

15           Properties of tablet composition :

pH of 20% (w/V) slurry : - Batch (J) = 5.92

- Batch (K) = 5.96

### **EXAMPLE 3 Amlodipine maleate tablets comprising mannitol**

20

#### **Composition**

Batch number	(L)
Equivalent of amlodipine base	10 mg
Amlodipine maleate	12.8 mg
Mannitol	370.1 mg
Sodium starch glycollate	8.0 mg
Magnesium Stearate	6.0 mg
Talc	2.0 mg
Total	398.9 mg

### Manufacturing procedure

- Amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate and talc have been mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- Magnesium stearate and talc were added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 10 mg tablets have been compressed using a Korsch EK0 excenter press.

### Example 4 Amlodipine maleate tablets having pH higher than 7 (comparative example)

Composition (pH of the 20% w/v slurry : 8.68 )

Batch number	(X)
Equivalent of amlodipine base	2.5 mg
Amlodipine maleate	3.21 mg
Calcium hydrogenphosphate anhydrous (pH 6.69)	31.5 mg
Magnesium oxide ponderosum	0.5 mg
Microcrystalline cellulose	62.05 mg
Sodium starch glycollate	2.0 mg
Magnesium Stearate	1.0 mg
Total	100.26 mg

### Manufacturing process

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- Amlodipine maleate, magnesium oxide and about 30 % of the amount of microcrystalline cellulose (MCC) were mixed in a free fall mixer for 10 minutes at about 25 rpm.
- The remaining amount of MCC, calcium hydrogenphosphate anhydrous and sodium starch glycollate were added and the blend was mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.

- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.

- 2.5 mg tablets and proportionately larger 10 mg tablets have been compressed using a Korsch EK0 excenter press.

5

#### **Example 5 Stability studies on amlodipine maleate tablets**

Stability studies on batches produced in Examples 1-4 were performed in a thermostated chamber adjusted to  $40 \pm 2$  °C and  $75 \pm 5$  % of relative humidity in various package materials ( HDPE bottles, PVC/PVDC/PE blisters ) or on an open dish. Assay of the active substance and of the content of impurities was performed by HPLC method, using reference materials of amlodipine maleate and major degradation impurities :

15 amlodipine aspartate ( Z#204)

amlo-pyridine (Z# 202)

Apart, two minor impurities Z#203 and Z#205 were detected and identified.

The content of other detected impurities/degradation products was calculated by internal normalization.

20 In the following tables, the assay of the active substance is expressed in milligrams, the content of impurities is expressed in per cent.

#### **A) Stability studies performed at 40°C/75%RH, open dish (influence of pH)**

<b>t=0 months</b>	<b>(A)</b>	<b>(B)</b>	<b>(E)</b>	<b>(F)</b>	<b>(C)</b>	<b>(D)</b>
Assay (mg/tab)	2.45	9.85	2.38	10.01	2.38	9.88
Z#202 (%)	0.15	0.15	0.08	0.08	0.05	0.08
Z#203 (%)	0.00	0.03	0.03	0.03	0.03	0.03
Z#204 (%)	0.16	0.15	0.13	0.12	0.13	0.12
Z#205 (%)	0.01	0.01	0.00	0.00	0.00	0.00
Tot.unknown(%)	0.31	0.28	0.27	0.28	0.27	0.27
<b>T= 1 month</b>						
Assay (mg/tab)	2.48	9.82	2.31	9.55	2.23	9.29
Z#202 (%)	0.15	0.15	0.21	0.23	1.27	1.25
Z#203 (%)	0.00	0.03	0.04	0.04	0.04	0.04
Z#204 (%)	0.24	0.23	0.41	0.30	0.27	0.22
Z#205 (%)	0.01	0.01	0.03	0.02	0.04	0.03
Tot.unknown(%)	0.35	0.31	0.58	0.56	0.83	0.72

<b>T= 3 months</b>						
Assay (mg/tab)	2.40	9.69	2.30	9.57	2.16	8.94
Z#202 (%)	0.20	0.18	0.37	0.39	1.85	1.85
Z#203 (%)	0.03	0.03	0.04	0.05	0.04	0.04
Z#204 (%)	0.31	0.29	0.77	0.63	0.33	0.27
Z#205 (%)	0.01	0.01	0.05	0.04	0.03	0.02
Tot. unkn. (%)	0.42	0.35	1.11	1.10	1.37	1.10

Stability studies performed in PVC/PE/PVDC blister at 40°C/75%RH

<b>T= 0 months</b>	<b>(A)</b>	<b>(B)</b>	<b>(J)</b>	<b>(K)</b>
Assay	2.51	9.99	2.52	10.33
Z#202	0.11	0.11	0.20	0.20
Z#203	0.00	0.00	0.00	0.00
Z#204	0.15	0.14	0.00	0.00
Z#205	0.01	0.01	0.00	0.00
t unkn.	0.31	0.31	0.46	0.49
<b>T= 3 months</b>				
Assay	2.49	9.49	2.49	9.99
Z#202	0.15	0.15	0.07	0.05
Z#203	0.00	0.00	0.00	0.03
Z#204	0.26	0.29	0.46	0.40
Z#205	0.01	0.01	0.02	0.02
t unkn.	0.36	0.34	0.45	0.38
<b>T= 6 months</b>				
Assay	2.45	9.49	2.45	9.94
Z#202	0.23	0.21	0.17	0.11
Z#203	0.04	0.04	0.04	0.04
Z#204	0.46	0.39	0.64	0.54
Z#205	0.01	0.00	0.02	0.02
t unkn.	0.57	0.54	0.64	0.51

5 Stability studies performed open dish at 40°C/75%RH

<b>t=0 months</b>	<b>(J)</b>	<b>(K)</b>
Assay	2.49	10.18
Z#202	0.04	0.04
Z#203	0.02	0.03
Z#204	0.19	0.17
Z#205	0.01	0.01
tot unkn.	0.42	0.36
<b>t=1 month</b>		
Assay	2.44	10.14
Z#202	0.08	0.07
Z#203	0.02	0.03
Z#204	0.34	0.36
Z#205	0.01	0.01
tot unkn.	0.45	0.42
<b>t=3 months</b>		

assay	2.41	9.72
Z#202	0.22	0.21
Z#203	0.04	0.04
Z#204	0.50	0.54
Z#205	0.03	0.03
tot unkn.	0.63	0.56

Stability studies performed in HDPE container at 40°C/75%RH

T= 0 months	(A)	(B)	(J)	(K)	(L)
Assay	2.51	9.99	2.52	10.33	10.11
Z#202	0.11	0.11	0.20*	0.20*	0.19*
Z#203	0.00	0.00	0.00	0.00	0.00
Z#204	0.15	0.14	0.00*	0.00*	0.00*
Z#205	0.01	0.01	0.00	0.00	0.00
t unkn.	0.31	0.31	0.46	0.49	0.40
T= 3 month					
Assay	2.52	9.84	2.51	10.26	8.69
Z#202	0.13	0.15	0.41*	0.37*	0.45*
Z#203	0.00	0.00	0.03	0.00	0.05
Z#204	0.27	0.23	0.00*	0.00*	0.00*
Z#205	0.03	0.03	0.07	0.05	0.01
t unkn.	0.33	0.32	0.41	0.49	0.66
T= 6 months					
Assay	2.49	9.83	2.47	10.26	
Z#202	0.15	0.15	0.08	0.05	
Z#203	0.00	0.00	0.00	0.03	
Z#204	0.44	0.41	0.44	0.38	
Z#205	0.02	0.01	0.10	0.08	
t unkn.	0.46	0.44	0.44	0.37	

\* Values for Z#202 and Z#204 together

5

Comparative stability studies performed at the composition with alkaline pH (open dish, 40°C/75%RH

t=0 months	(X)
Assay	2.52
Z#202	0.04
Z#203	0.03
Z#204	0.13
Z#205	0.00
tot unkn.	0.30
t=1 month	
Assay	2.43
Z#202	0.06
Z#203	0.03
Z#204	1.73
Z#205	0.00

tot unkn.	0.51
-----------	------

Stability studies performed at 40°C/75%RH, open dish (influence of particle size)

<b>t = 0 months</b>	(A)	(B)	(G)	(H)	(I)
Assay (mg/tab)	2.45	9.85	2.38	10.12	2.54
Z#202 (%)	0.15	0.15	0.05	0.05	0.10
Z#203 (%)	0.00	0.03	0.00	0.01	0.04
Z#204 (%)	0.16	0.15	0.03	0.03	0.01
Z#205 (%)	0.01	0.01	0.00	0.00	0.02
Tot unkn. (%)	0.31	0.28	0.17	0.11	0.26
<b>t = 1 month</b>					
Assay (mg/tab)	2.48	9.82	2.29	9.88	-
Z#202 (%)	0.15	0.15	0.17	0.14	-
Z#203 (%)	0.00	0.03	0.01	0.01	-
Z#204 (%)	0.24	0.23	0.52	0.48	-
Z#205 (%)	0.01	0.01	0.02	0.02	-
Tot unkn. (%)	0.35	0.31	0.22	0.19	-
<b>t = 2 months</b>					
Assay (mg/tab)	2.46	9.87	2.22	9.64	2.35
Z#202 (%)	0.17	0.16	0.28	0.25	0.34
Z#203 (%)	0.00	0.00	0.01	0.01	0.03
Z#204 (%)	0.28	0.28	1.02	1.01	0.41
Z#205 (%)	0.01	0.01	0.03	0.03	0.04
Tot unkn. (%)	0.47	0.43	0.63	0.52	0.60
<b>t = 3 months</b>					
Assay (mg/tab)	2.40	9.69	2.24	9.36	2.36
Z#202 (%)	0.20	0.18	0.18	0.16	0.39
Z#203 (%)	0.03	0.03	0.01	0.01	0.11
Z#204 (%)	0.31	0.29	1.26	1.34	0.53
Z#205 (%)	0.01	0.01	0.05	0.04	0.04
Tot unkn. (%)	0.42	0.35	0.90	0.74	0.64

5

For a comparison of stability, increase in total impurities, identified and unidentified, with regard to the t=0 months level, is considered:

increase in total impurities in %	(A)	(B)	(G)	(H)	(I)
after 3 months at 40°C/75%RH	+ 0.33	+ 0.24	+ 1.82	+ 1.43	+1.28

10

**Comparative stability study with Norvasc (commercial amlodipine besylate tablets)**

Stability studies performed in original blister at 40°C/75%RH

	Norvasc® 5 mg batch 81040100 (DE)	Norvasc® 10 mg batch 901-05941 (NL)
<b>t = 0 months</b>		
Assay	5.19	9.99
Z#202	0.04	0.02
Z#203	0.01	0.00
Z#204	0.00	0.00
Z#205	0.00	0.00
tot. unkn.	0.12	0.12
<b>t = 3 months</b>		
Assay	5.13	9.70
Z#202	0.16	0.17
Z#203	0.03	0.00
Z#204	0.00	0.00
Z#205	0.00	0.00
tot. unkn.	0.38	0.62
<b>t = 6 months</b>		
Assay	4.97	9.58
Z#202	0.28	0.27
Z#203	0.00	0.00
Z#204	0.00	0.00
Z#205	0.00	0.00
tot. unkn.	0.49	0.78

Stability studies performed on open dish at 40°C/75%RH

		Norvasc® 2.5 mg batch 8QP115A (US)	Norvasc® 10 mg batch N-09 (ES)
<b>t = 0 months</b>	Assay	2.44	9.91
	Z#202	0.08	0.82
	Z#203	0.00	0.00
	Z#204	0.00	0.00
	Z#205	0.00	0.00
	tot. unkn.	0.02	0.34
<b>t = 1 months</b>	Assay	2.44	8.90
	Z#202	0.18	2.17
	Z#203	0.04	0.19
	Z#204	0.00	0.01
	Z#205	0.00	0.00
	tot. unkn.	0.10	1.21
<b>t = 2 months</b>	Assay	2.39	7.98
	Z#202	0.27	3.24
	Z#203	0.00	0.00
	Z#204	0.00	0.00
	Z#205	0.00	0.03
	tot. unkn.	0.33	2.51
<b>t = 3 months</b>	Assay	2.34	7.68
	Z#202	0.37	3.98
	Z#203	0.00	0.00

	Z#204	0.00	0.00
	Z#205	0.00	0.03
	tot. unkn.	0.27	2.76

#### **Example 6 Amlodipine maleate capsules**

5

#### **Compositions**

Batch number	(CA), (CB)
Equivalent to amlodipine	5.0 mg
Amlodipine maleate	6.42 mg
Microcrystalline cellulose	72.6 mg
Predried potato starch	20.0 mg
Magnesium stearate	0.5 mg
Total	99.52 mg

Batch number	(CC)	(CX)
Equivalent to amlodipine	5.0 mg	5.0 mg
Amlodipine maleate	6.42 mg	6.42 mg
Calcium hydrogenphosphate anhydrous	31.5 mg	31.5 mg
Magnesium oxide	-	1.5 mg
Microcrystalline cellulose	62.0 mg	62.0 mg
Sodium starch glycollate	2.0 mg	2.0 mg
Magnesium stearate	1.0 mg	1.0 mg
Total	102.92 mg	104.43 mg

10

pH of capsules at t=0

Batch no.	Strength	Type of Calcium hydrogenphosphate anhydrous	pH of 20% (m/V) slurry of capsule content
(CC)	5.0 mg	Di CAFOS A	6.10
(CX)	5.0 mg	Di CAFOS A	9.59

#### **Batch (CA) has been manufactured as follows:**

15

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.

- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.

- *gelatine* capsules have been filled with this powder blend

5        **Batch (CB) has been manufactured as follows:**

- The amlodipine maleate was sieved through a 500 µm screen.

- The other excipients have been sieved through a 850 µm screen.

- All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.

10        - Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.

- *HPMC* capsules have been filled with this powder blend

15        **Batch (CC) has been manufactured as follows:**

- The amlodipine maleate was sieved through a 500 µm screen.

- The other excipients have been sieved through a 850 µm screen.

- All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.

20        - Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.

- *gelatine* capsules have been filled with this powder blend

25        **Batch (CX) has been manufactured as follows:**

- The amlodipine maleate was sieved through a 500 µm screen.

- The other excipients have been sieved through a 850 µm screen.

- Amlodipine maleate, magnesium oxide and about 30 % of the amount of microcrystalline cellulose (MCC) were mixed in a free fall mixer for 10 minutes at about 25 rpm.

30        - The remaining amount of MCC, calcium hydrogenphosphate anhydrous and sodium starch glycollate were added and the blend was mixed in a free

fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.

- Magnesium stearate was added and the powderblend was mixed for another 5 minutes at about 25 rpm.

- 5        - *gelatine* capsules have been filled with this powderblend, using an automatic capsule filling machine.

### **Example 7 Stability studies on amlodipine maleate capsules**

- 10        Stability studies on batches prepared in Example 6 were performed essentially as described in Example 5.

Stability studies performed in PVC/PE/PVDC blister at 40 °C/75 % RH

<b>t=0 months</b>	<b>(CA)</b>	<b>(CB)</b>
Assay	5.13	4.98
Z#202	0.04	0.04
Z#203	0.03	0.03
Z#204	0.12	0.12
Z#205	0.00	0.00
tot unkn.	0.32	0.31
<b>t=3 months</b>		
Assay	4.82	4.76
Z#202	0.08	0.06
Z#203	0.02	0.02
Z#204	0.15	0.14
Z#205	0.01	0.00
tot unkn.	0.39	0.38
<b>t=6 months</b>		
Assay	4.67	4.81
Z#202	0.14	0.10
Z#203	0.04	0.02
Z#204	0.23	0.18
Z#205	0.02	0.01
tot unkn.	0.45	0.42

15

Two one month stability studies performed in open dish

<b>t=0 months</b>	<b>(CC)</b>	<b>(CX)</b>
Assay	4.91	4.72
Z#202	0.04	0.04

Z#203	0.03	0.04
Z#204	0.12	0.13
Z#205	0.00	0.00
tot unkn.	0.28	0.29
T = 1 month @ 25°C/60%RH		
Assay	4.85	4.70
Z#202	0.05	0.05
Z#203	0.04	0.04
Z#204	0.13	0.14
Z#205	0.00	0.00
tot unkn.	0.28	0.28
T=1 month @ 40°C/75%RH		
Assay	4.76	4.16
Z#202	0.06	0.08
Z#203	0.04	0.03
Z#204	0.15	11.36
Z#205	0.00	0.00
tot unkn.	0.28	0.62

Stability studies performed in HDPE container

<b>t=0 months</b>	<b>(CA)</b>	<b>(CB)</b>
Assay	4.99	5.01
Z#202	0.05	0.04
Z#203	0.03	0.03
Z#204	0.12	0.11
Z#205	0.00	0.00
tot unkn.	0.31	0.30
T= 3 months @ 25°C/60%RH		
Assay	5.07	4.77
Z#202	0.04	0.04
Z#203	0.03	0.04
Z#204	0.12	0.12
Z#205	0.00	0.00
tot unkn.	0.30	0.32
T= 3 months @ 40°C/75%RH		
Assay	5.01	4.65
Z#202	0.05	0.05
Z#203	0.03	0.04
Z#204	0.12	0.13
Z#205	0.00	0.00

tot unkn.	0.29	0.32
-----------	------	------

Stability studies on commercially available Amlor® (amlodipine besylate) capsules

- 5 Stability studies performed in original blister

<b>Amlor 5 mg capsules batch 9037002</b>	<b>t=0 months</b>	<b>t=3 months 40°C/75%RH</b>
Assay	4.59	4.44
Z#202	0.01	0.20
Z#203	0.00	0.00
Z#204	0.00	0.00
Z#205	0.00	0.00
tot unkn.	0.06	0.37

- 10 The invention having been described, it will be readily apparent to those skilled in the art that further changes and modifications in actual implementation of the concepts and embodiments described herein can easily be made or may be learned by practice of the invention, without departing from the spirit and scope of the invention as defined by the following claims